

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Dedicate®

TURF AND ORNAMENTAL FUNGICIDE

**ACTIVE CONSTITUENTS: 200 g/L TEBUCONAZOLE
100 g/L TRIFLOXYSTROBIN**

GROUP 3 11 FUNGICIDE

* For the control of fungal diseases in turf and ornamentals as per the Directions for Use.

* 500 mL – 3 L

DIRECTIONS FOR USE

RESTRAINTS

SPRAY DRIFT RESTRAINTS

DO NOT apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturer specifications that refer to ASAE S572 Standard to the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are:

1. date and start and finish times of application;
 2. location address and areas sprayed;
 3. full name of this product;
 4. amount used per hectare and number of hectares applied to;
 5. situation and disease;
 6. wind speed and direction during application;
 7. air temperature;
 8. nozzle brand, model and type and spray system pressure measured during application;
 9. name and address of person applying this product.
- (Additional record details may be required by the State or Territory where this product is used).

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas, including aquacultural ponds or surface streams and rivers or, within **10 metres** downwind from the application area.

DO NOT apply if there are susceptible non-target plants within **15 metres**, downwind from the application area.

DO NOT apply if there are livestock, pasture or any land that is producing feed for livestock within **15 metres** downwind from the application area.

DIRECTIONS FOR USE

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Turf	Anthracnose (<i>Colletotrichum graminicola</i>); Brown Patch (<i>Rhizoctonia solani</i>); Fusarium or Microdochium Patch (<i>Microdochium nivale</i>); Helminthosporium disease (<i>Bipolaris</i> spp., <i>Drechslera</i> spp., <i>Exserohilum</i> spp.); Leptosphaerulina; Curvularia	2 L/ha in 400 – 1600 L water per ha or 20 mL/100 m ² in 4 – 16 L water per 100 m ²	For preventative control apply on a 21-28 day interval when conditions are favourable for disease development. For curative control apply on a 14 – 21 day interval. When targeting brown patch control on the roots, apply sufficient irrigation to wash Dedicate into the root zone. This use is subject to an FRAC anti- resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.
	Ectotrophic Root Infecting Fungi (ERI)		Make monthly preventative applications in the three months preceding the disease season. Apply irrigation to wash Dedicate into the root zone. This use is subject to an FRAC anti- resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.
	Spring Dead Spot (<i>Ophiosphaerella namari</i>)		Apply as a preventative application in January to March after renovation and recovery of active growth. Make a second application one month later. Apply irrigation to wash Dedicate into the root zone.
	Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	3 L/ha in 400 – 1600 L water per ha	For preventative control apply on a 21-28 day interval when conditions are favourable for disease development. To control existing infections apply on a 14 – 21 day interval. This use is subject to an FRAC anti- resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.
	White helminthosporium	280 mL per bowling green in 150 L water per bowling green	Make monthly preventative applications from December to March. This use is subject to an FRAC anti- resistance strategy. Do not apply more than two consecutive sprays of Dedicate unless mixed with a protectant fungicide from group 14, 28, M2, M3, M4 or M5.

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Azaleas	Azalea petal blight (<i>Ovulinia azaleae</i>)	75 mL / 100 L water	Apply when disease symptoms first appear and then repeat at 7 – 10 day intervals. When conditions favour disease development the shorter spray interval should be considered. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves.
Ornamentals	Botrytis blight or Grey mould (<i>Botrytis cinerea</i>); Downy Mildew (<i>Peronospora</i> spp.); Leaf spots (<i>Pseudomonas</i> spp.); Rusts (<i>Uromyces erythronii</i> ; <i>Puccinia</i> spp.)	75 mL / 100 L water	Curative: For existing infections, spray to run-off ensuring thorough coverage of all foliage including the underside of leaves in a series of 3 applications at 5 day intervals. Preventive: Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 7 to 14 days according to infection pressure or beginning immediately after onset of the disease.
Roses	Black Spot (<i>Diplocarpon rosae</i>)		
Ornamentals; Roses	Powdery mildew (<i>Sphaerotheca pannosa</i> var. <i>rosae</i> ; <i>Erysiphe cichoracearum</i> ; <i>Oidium</i> spp.)	150 mL / 100 L water	Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 7 - 14 days according to infection pressure or beginning immediately after onset of the disease.
Ornamentals and non-fruit- bearing plants of the Myrtaceae family	Myrtle Rust (<i>Uredo rangalii</i>)	75 mL / 100 L water	Apply at first signs of disease or when conditions favour disease development. Curative: For existing infections, apply 2 applications at a 14-day interval. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. Preventive: Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 14 days according to infection pressure.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIOD

DO NOT graze treated turf; or feed turf clippings from any treated area to poultry or livestock.

GENERAL INSTRUCTIONS

APPLICATION

Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage. For application in turf, use 400-800 L water per ha for foliar or crown diseases such as Dollar Spot, Anthracnose, Brown Patch, Fusarium (Microdochium), Helminthosporium, Leptosphaerulina and Curvularia. Use higher water rates (up to 1600 L per ha) for treatment of soil/root diseases such as Brown Patch, Ectotrophic Root Infecting fungi and Spring Dead Spot.

Dedicate Turf and Ornamental Fungicide has been used on a range of ornamental plant species (i.e. azaleas, begonias, chrysanthemums, cinerea, geraniums, gerbera, Myrtaceae, native daisy, pelargoniums, roses, snapdragon) without damage. Some species and varieties however are particularly sensitive to chemical sprays and as this sensitivity is often related to local conditions, it is advisable to treat only a small number of plants first, in order to ascertain the reaction before treating the whole target group.

TURF NUTRITION

This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertiliser application. Where possible, it is recommended that the fertiliser program should precede the fungicide application.

IPM COMPATIBILITY

RISK TO NON-TARGET INSECTS: DEDICATE may have adverse effects on some nontarget beneficial insects, in particular predatory bugs. This risk should be taken into account where Integrated Pest Management utilising these organisms is practised.

RESISTANCE WARNING

GROUP	3	11	FUNGICIDE
-------	---	----	-----------

For fungicide resistance management DEDICATE is a Group 3 & 11 fungicide. Some naturally occurring individual fungi resistant to DEDICATE and other Group 3 & 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by DEDICATE and other Group 3 & 11 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, 2022 Environmental Science AU Pty Ltd accepts no liability for any losses that may result from the failure of DEDICATE to control resistant fungi.

PRECAUTIONS

RE-ENTRY PERIOD FOR TURF

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

RE-ENTRY PERIOD FOR ORNAMENTALS

DO NOT enter treated areas until spray has dried for low and medium exposure activities such as irrigation, scouting, thinning and weeding of immature/low foliage plants and/or mature/high foliage plants, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves. Clothing must be laundered after each day's use.

DO NOT enter for five (5) days after application for high exposure activities such as hand harvesting, pruning and pinching, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS

When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing buttoned to the neck and wrist and elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (ph: 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which may be found at www.au.envu.com.

CONDITIONS OF SUPPLY

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE RETURN AT ONCE UNOPENED.

To the extent permitted by law 2022 Environmental Science AU Pty Ltd does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. 2022 Environmental Science AU Pty Ltd does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter. To the extent only that 2022 Environmental Science AU Pty Ltd is prevented by law from contracting out of the Consumer Guarantees Act then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

2022 Environmental Science AU Pty Ltd
ABN 49 656 513 923
Suite 2.06, Level 2, 737 Burwood Road,
Hawthorn East, Vic. 3123
Technical enquiries 1800 024 209
technicalsupport.australia@envu.com
www.au.envu.com

Envu, the Envu logo and DEDICATE® are trademarks owned by Environmental Science U.S. LLC or one of its affiliates.

©2024 Environmental Science U.S. LLC

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY CALL +61 2 9037 2994 or 1800 862 115 toll free. The toll free phone number is possibly accessible, but not guaranteed from payphones within Australia and is not accessible from outside of Australia.

BN:
DOM:



GHS statement

Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. Very toxic to aquatic life with long lasting effects.

Do not handle until all safety precautions have been read and understood. Do not breathe mist. Avoid contact during pregnancy/while nursing. Do not eat, drink or smoke when using this product. Use personal protective equipment as required. If exposed or concerned: get medical advice/attention. Store locked up.