

# POISON

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY INSTRUCTIONS BEFORE  
OPENING OR USING**



**Oxa-Pro® 15:2:8**

**FERTILISER WITH PRE-EMERGENT HERBICIDE**

**ACTIVE CONSTITUENTS: 10g/kg OXADIAZON  
SOLVENT: 10.5g/kg POLYBUTENE**

**GROUP G HERBICIDE**

**GUARANTEED FERTILISER ANALYSIS**

15.0%	TOTAL (N)
2.0%	As ammoniacal N
13.0%*	As urea N
2.2%	PHOSPHORUS (P)
8.3%	POTASSIUM (K)
2.4%	SULPHUR (S) as elemental

\*6.0% Slowly Available Nitrogen from Polymer Coated Sulphur Coated Urea

**COMBINATION HERBICIDE AND CONTROLLED RELEASE FERTILISER  
FOR PRE EMERGENT CONTROL OF SUMMER GRASS & CRAB GRASS  
AND WINTER GRASS IN TURF and FEEDING OF TURF WITH SLOW  
RELEASE NITROGEN WITH PHOSPHORUS AND POTASSIUM**

**NET CONTENT: 20 kg**



**BARMAC INDUSTRIES PTY LTD  
3 Mary Street  
Blackstone, QLD 4304**

APVMA Approval number 59774/20kg/0805

**DIRECTIONS FOR USE**

**Restraints: DO NOT** apply granules when the turf foliage is wet.  
**DO NOT** use on putting greens, low cut tees or bowling greens.  
**DO NOT** apply to newly seeded turf areas.

Situation	Target	Turf Species	Rate	Critical Comments
Warm season turfgrass areas, including fairways, golf tees, municipal sports fields, industrial and commercial lawns	Pre emergent control of Summer Grass ( <i>Digitaria ciliaris</i> ) Crowsfoot Grass ( <i>Elcusine indica</i> )	Kikuyu ( <i>Pennisetum clandestinum</i> ), Buffalo Grass ( <i>Stenotaphrum secundatum</i> ), Common Couch ( <i>Cynodon dactylon</i> , <i>C. transvaalensis</i> ) Grass (Not Santa Ana)	300 kg/ha (3kg/100m <sup>2</sup> )	The product is ready to apply. No mixing or dilution is required. Apply product evenly to dry foliage and bare areas using a granular spreader able to give even coverage. Application must be made before weeds have begun to germinate. Established weeds will not be controlled. After application water in with at least 10mm of sprinkler irrigation. Control of germinating weed seedlings can be expected for up to 10 weeks after application depending upon temperature, rainfall, soil type and other factors. If longer control is required a repeat application should be made 7-10 weeks after initial application. <b>DO NOT</b> mow treated area until product has been watered in and turf foliage has dried. Avoid application when daytime temperatures are consistently above 32°C. This product is not recommended for use on putting greens, low cut tees or bowling greens. <b>DO NOT</b> apply on turf containing Carpet Grass ( <i>Axonopus</i> spp.), Centipede Grass ( <i>Eremochloa ophiuroides</i> ) and Dichondra unless some damage is acceptable.
	Pre emergent Control of Winter Grass ( <i>Poa annua</i> ) Creeping Oxalis ( <i>Oxalis comiculala</i> )	Tif Hybrid Couch Grass, Saltwater couch ( <i>Paspalum vaginatum</i> ), Queensland Blue Couch ( <i>Digitaria didactyla</i> ), Bahla grass ( <i>Paspalum notatum</i> ), Zoysia grass ( <i>Zoysia matrella</i> )	400kg/ha (4kg/100m <sup>2</sup> )	
	Turf nutrition - provides slow release nitrogen for 8 weeks feeding		300-400kg/ha (3-4kg/100m <sup>2</sup> )	

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**GENERAL INSTRUCTIONS:**

The Andersons Oxa-Pro® 15:2:8 Fertiliser with Pre-Emergent Herbicide is a convenient ready to use formulation for the control of certain annual grass weeds in turf and also for fertilisation of turf where nitrogen, potassium and sulphur are required. The product should only be used when both functions are required.

**APPLICATION:**

For a granular spreader which can be calibrated for even output and capable of giving even application. Use a double application technique: i.e. half rate width, to achieve a full rate application. Apply under calm conditions, thereby reducing drift to non-target areas and optimizing particle distribution for good weed control.

Ensure turf area is free of leaves and other debris, mowing if necessary so that the granules reach the soil surface. Delay reseeding of Ryegrass or Bentgrass into treated areas for 4-5 months (Ryegrass 4 months & Bentgrass 5 months). **DO NOT** over apply, especially under the branch spread of trees and shrubs. Avoid contact with flowers and shrubs.

Suggested Spreader Settings				
The suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.				
Spreader	Ground Speed	Width of Coverage	SPREADER SETTINGS	
			300 kg/ha	400 kg/ha
Scotts AccuPro 2000, SR-2000 (cone 4)	4.8 km/hr	4.0 m	O 1/2	Q
Scotts R-8, R-8A (cone 4)	4.8 km/hr	4.0 m	O	P 1/2
Spyker 288 Super	4.8 km/hr	3.0 m	3.9	4.2
Vicon (03 Series - Standard Spout)	7.3 km/hr	11.0 m	37	44

**HERBICIDE RESISTANCE WARNING:**

**GROUP G HERBICIDE**

Oxadiazon Herbicide is a member of the Oxadiazole Group of herbicide. The herbicide is an inhibitor of protoporphyrinogen oxidase. For weed resistance management, the product is a Group G Herbicide. Some naturally occurring weed biotypes resistant to the herbicide may exist through normal genetic variability in weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other oxadiazole group herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Barmac Industries Pty Ltd does not accept liability for any losses that may result from failure of this product to control resistant weeds.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:**

Dangerous to Fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply product if rapid storm run-off from treated areas into water bodies is likely to occur within several days of application. **STORAGE AND DISPOSAL:** Store in the closed original container, in a dry, cool well ventilated area out of direct sunlight. Do not store with seed, feed, foodstuffs, insecticides, fungicides and other fertilisers. Shake bag empty into applicator. Puncture and shred empty container and bury in a local authority landfill. If no landfill is available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. DO NOT dispose of damaged material on site. DO NOT re-use this container for any purpose.

**SPILLS:** Sweep up spilled materials and return to original container. Wash contaminated equipment taking care the washings DO NOT enter sewers, drains and other waterways.

**SAFETY DIRECTIONS:** Avoid contact with eyes. Do not initiate dust. Wash hands after use.

**FIRST AID:** If poisoning occurs contact a doctor or Poisons Information Centre Australia 131126.

**MATERIAL SAFETY DATA SHEET:** Further information on this product is contained in a MSDS, copies of which can be obtained from our web site, [www.barmac.com.au](http://www.barmac.com.au)

**NOTICE:** This product is designed only for the particular purpose indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Barmac Industries Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or other wise in connection with the supply and use for this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.

Batch Number \_\_\_\_\_  
Date of Manufacture \_\_\_\_\_

F7b (APPBLKPDNAB)

**APPBMC8684-1**

**FAN 8684-1**