

# CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Trimmit<sup>®</sup>

Growth Regulator

syngenta<sup>®</sup>

**ACTIVE CONSTITUENT: 250 g/L PACLOBUTRAZOL**

Controls growth in hedging plants as specified in the  
Directions for Use

**Syngenta Australia Pty Ltd**

Level 1, 2-4 Lyonpark Road, Macquarie Park NSW 2113

***In a transport emergency dial 000, Police or Fire Brigade***

***For specialist advice in an emergency only, call 1800 033 111 (24 hours)***

**APVMA Approval No: 82071/104922**

TM

## DIRECTIONS FOR USE

### Restraints

DO NOT use on plants of low vigour or under stress (see General Instructions)

DO NOT apply to newly planted or recently transplanted hedging species, allow hedge to establish and consolidate before applying for the first time

Situation	State	Application	Rate	Critical Comments
<b>Hedging species</b> Golden Duranta <i>(Duranta repens)</i> , Small leaf Photinia <i>(Photinia glabra)</i> , Syzygium <i>(Syzygium australe, Syzygium smithii)</i>	All States	Root drench	1.5 to 2.5 mL in 1 L water/plant	Make applications to the hedging plants in late winter to early spring, ideally at the commencement of a new season flush of growth. Applications must cover the root system adjacent to the stem at ground level. Remove mulch to ensure application reaches soil. Light post application irrigation is recommended. DO NOT apply to hedges less than 0.3 m or greater than 2 m in height It is critical the apical growing points of the hedging plant be in tact to act as sinks drawing the product into the plant at the time of application. Therefore, DO NOT prune the hedging plant until 4 to 6 weeks after application. DO NOT allow plants to experience drought stress just prior to or just after application. DO NOT make a repeat treatment within the following 12 months. DO NOT apply to newly planted or recently transplanted hedging species. It is strongly recommended the lower label rates be used when applying for the first time. The higher rate can be used in following years when the responsiveness of the hedge is established.
<b>Hedging species</b> Mock Orange <i>(Murraya paniculata)</i>			0.5 to 1.5 mL in 1 L water/plant	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

### GENERAL INSTRUCTIONS

TRIMMIT is a systemic growth regulator that reduces internode lengths of new shoots and causes earlier formation of terminal buds. Use of TRIMMIT should begin only when plants are nearing their optimum size. For hedges this is once the desired shape (height, width and foliar density) has already been achieved and the hedge requires only maintenance thereafter. Responsiveness to this product can vary from site to site. It is strongly recommended the lower rate be used when applying for the first time. If increased regulation is required then step up to the higher rate in the second year. If a plant is under stress, TRIMMIT should NOT be used. Stress can be caused by moisture shortage due to lack of rain or inadequate irrigation, high temperature, salinity, moisture excess due to water table, flooding or inappropriate irrigation, severe or successive frosts, poor fertility caused by imbalance of major or minor nutrients, diseases, insects, inappropriate herbicide use, mechanical damage.

### Hedge Management

Treated hedges should be kept well watered with adequate nutrition. Plants should not have nutrition limited due to a lesser growth rate. When replanting a hedge at least 2 years should have elapsed since the last application of TRIMMIT. Alternatively, if planting into an area that historically has been treated, fresh soil should be provided to allow plant establishment and growth. A further alternative is to avoid original planting holes and place young hedging species as far as practicable from the treated soil (minimum 1.5 m).

## Mixing

Mix required volume of TRIMMIT with recommended volume of water. Maintain agitation to prevent settling.

## Application

**Root Drench:** Uptake of TRIMMIT is most efficient when it is applied to soil which receives adequate water and where there is a high density of feeder roots. For hedging species apply required amount of TRIMMIT in 1 L water per plant. Ensure all treated plants are lightly irrigated following treatment under dry conditions. Individual trees or plants which are smaller than average should be left untreated or given a proportionally smaller dose of TRIMMIT.

## Retreatment

In some cases, control of growth may persist for more than 1 year. Retreat plants only when normal growth resumes. Only retreat in the late winter or spring that follows resumption of normal growth. Use the interval between the first treatment and the resumption of normal growth as a guide for retreatment in subsequent seasons.

## Compatibility

The compatibility of TRIMMIT with other products has not been established. Always conduct a physical compatibility test and check for plant safety on a sample plant before making broader applications of mixes to hedging species.

## PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

## SAFETY DIRECTIONS

**Avoid contact with the eyes and skin. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.**

## FIRST AID

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.**

## MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 067 108, or visit our website at [www.greencast.com.au](http://www.greencast.com.au) or [www.syngenta.com.au](http://www.syngenta.com.au)

## DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company

