



HERBICIDE REFERENCE GUIDE

Product	Active(s)	Group	Rate / 100m ²	Water Volume (L / 100m ²)	Pre / Post Emergent	Broadleaf											Grass Weeds						Sedges		Non-selective		Landscape Weeds		Post Application Requirements	Adjuvants recommended?								
						Algae	Bindii	Burr Medic	Capeweed	Clover	Cudweed	Fleabane	Chickweed	Catsear	Creeping Oxalis	Plantain	Pearlwort	Wireweed	Rye Grass	Summer Grass	Winter Grass	Couch	Paspalum	Kikuyu	Crowsfoot	Mullumbimby Couch	Nut Grass	Onion Grass			Various	Various						
						Annual					Perennial						Annual		Perennial		Both		Perennial		Both													
3D	Bentazone, MCPA & Dicamba	C1	200mL	20	Post																																Apply to young, vigorous annuals and prior to flowering for perennials	Yes
Algaefix	Q.A.C	-	50mL	5	Post																															Leave on foliage. Apply at 7-10 day intervals, as required		
Barricade	Proflamifone	D	10 - 30mL	> 5	Pre																															Apply to foliage in advance if germination for up to 6 months residual	Yes	
Basta	Glufosinate - ammonium	N	10 - 50mL	10	Post																															Apply to foliage when targets are actively growing and not under stress	Yes	
Bronco MA-X	Bromoxynil & MCPA	C I	43mL	15	Post																															Apply to foliage	Yes	
Casper	Dicamba & Prosulfuron	B I	6 - 12g	4 - 8	Post																															WDG product. Quick systemic movement	Yes	
Coliseum	Rimsulfuron	B	0.75 - 1.2g	4	Post																															Apply to foliage	Yes	
Crest	Florasulam & Fluroxypyr	B I	10 - 20mL	0.8 - 1	Post																															Apply to foliage	Yes	
Crowbar	Diclofop - methyl	A	10 - 25mL	> 5	Post																															Apply to foliage	Yes	
Destiny	Iodosulfuron - methyl	B	1 - 1.5g	2 - 5	Post																															Apply to foliage	Yes	
Dimension EW	Dithiopyr	D	17.5 - 35mL	2 - 4	Pre & Post																															Apply to foliage	Yes	
DithoMAX	Dithiopyr	D	2.5Kg	N / A	Pre																															19:0:5 fertiliser prill with Pre emergent herbicide coating		
Double Time	Dicamba & MCPA	I	65mL	2.4 - 4.7	Post																															Apply to foliage	Yes	
Drive XL	Quinclorac	I	46mL	4	Post																															Apply to foliage, ideally in Autumn	Yes	
DSMA Clear	DSMA	K	250 - 660mL	4.5 - 20	Post																															Apply to foliage	Yes	
Embargo	Oryzalin	D	50mL	4 - 8	Pre																															Follow up application with 10-15mm of irrigation within 24 hours	Yes	
Eraze 360	Glyphosate	M	12.5 - 30mL	2 - 4	Post																															Non selective	Yes	
Eraze 510	Glyphosate	M	10 - 60mL	2 - 4	Post																															Non selective	Yes	
Jolt / Spearhead	MCPA, Clopyralid & Diflufenican	F I	50mL	2 - 5	Post																															Apply to foliage of actively growing weeds	Yes	
Matrix	Ethofumesate	J	8mL	4 - 6	Pre & Post																															Apply one month before expected germination. Dont Irrigate for 12 hours after application.	Yes	
Meteor	Metolachlor	K	10 - 20mL	4	Pre																															Wash in with 2 - 4mm of irrigation	Yes	
Metric	Metribuzin	C	10mL	4 - 8	Early Post																															Leave on the foliage of emerged weeds.	Yes	
Mojo	Sulfometuron - methyl	B	2 - 4g	> 0.5	Post																															WDG.	Yes	
Monopoly	MSMA	Z	66mL	4 - 10	Post																															Split application at 7 days if phytotoxicity risk is undesirable	Yes	
Monument	Trifloxysulfuron Sodium	B	1.15 - 3mL	4 - 8	Post																															Leave on foliage - ideal application is in Autumn	Yes	
Nominee	Bisoyribac - Sodium	B	10mL	> 2.5	Post																															2 applications spaced 2 - 4 weeks apart is suggested	Yes	
Oxa MAX (18:10:9)	Oxadiazon	G	3 - 4Kg	N / A	Pre																															Follow application with at least 10mm of irrigation		
Pac Down	Paclbutrazol	PGR	16 - 22mL	4 - 8	Post																															Use 4 - 6mm of irrigation immediately after application. Only apply to healthy turf. Supression of Poa only.	Yes	
Pendant 330	Pendimethalin	D	30 - 45mL	> 4	Pre																															Leave on foliage - ideally programmed spring applications are applied	Yes	
PendiMAX (22:0:5)	Pendimethalin	D	1.6 - 2.6Kg	N / A	Pre																															Irrigate with 10mm after application to activate nutrition and pre emergent		
Pronamide	Propyzamide	D	6 - 12mL	2 - 6	Pre and early Post																															Apply during winter and Autumn. Immediately irrigate with 12 - 25mm and again 24 hrs later	Yes	
Ronstar G	Oxadiazon	G	1.5 - 2Kg	N / A	Pre																															Granular. Apply through irrigation if rainfall hasn't fallen within 48 hrs.		
Sedgehammer	Halosulfuron - methyl	B	0.65 - 1.3g	10	Post																															Apply to the foliage of healthy, actively growing selected weeds	Yes	
Silverado	Carfentrazone & MCPA	G I	10g	8 - 16	Post																															Works well on Moss. Repeat applications after 14 days is encouraged.	Yes	
Smackdown	Carfentrazone - ethyl	G	0.25 - 0.75g	3 - 6	Post																															Ideal supplementary partner for non selective herbicide (Glyphosate)	Yes	
Specticle	Indaziflam	O	2.5mL	2 - 5	Pre																															Good residual. Apply prior to weed germination to healthy turf	Yes	
Steer 750	Quinclorac	I	11g	4	Post																															Kikuyu supression only. Apply to foliage	Yes	
Stonewall	Oryzalin	D	45 - 68mL	2 - 4.5	Pre																															Leave on foliage in advance of weed emergence	Yes	
Tribute	Foramsulfuron	B	15 - 20mL	> 2	Post																															Kikuyu supression only. Apply to foliage. Some treatments require repat application.	Yes	
Triple Shot / Triple Time	MCPA, Bromoxynil & Dicamba	C I	5mL	2.4 - 4.7	Post																															Leave on foliage	Yes	
Tupersan	Siduron	C	1Kg	10 - 20	Post																															Wettable Powder. Apply at beginning and end of cooler season for warm season grassy weed control	Yes	
Turzine Pro	Isoxaben & Florasulam	O B	0.75 - 1g	4	Pre & Post																															WDG. Leave on foliage of actively gorwing weeds.	Yes	
Unimaz	Imazapyr	B	40mL	0.5 - 2	Pre & Post																															Leave on foliage	Yes	
Vault	Oxyfluorfen & Oryzalin	DG	1Kg	2 - 5	Pre																															Leave on foliage and follow up with 10-15mm of irrigation	Yes	



Nuturf's Technical Department provides a fast, accurate disease diagnostic service.

The report includes recommendations both from a cultural and chemical control perspective.

For more information on this service, enquire with your Nuturf Territory Manager.



Contact your Nuturf Territory Manager for more information
 1800 631 008 | www.nuturf.com.au