

**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PEEL HERE



**Pride**

Insecticide

ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP

**4A**

INSECTICIDE

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT:**  
**READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**

Amgrow Pty Ltd (Commercial Products Division)

ABN 22 069 900 456

Unit B2A, 3-29 Birnie Ave,

Lidcombe NSW 2141

TEL: 1800 631 008

[www.ecofertiliser.com.au](http://www.ecofertiliser.com.au)



**CONTENTS**

**1L**

© Pride is a registered trademark of Ecofertiliser Pty Ltd.

APVMA Approval No. 64900/1L/0410



File Name:

64900\_49555\_1L\_label\_MPL\_V2.pdf

Dimensions: 100mm wide x 160mm high

Scale of Print: A4 100% A3 100%

## DIRECTIONS FOR USE

CROP	PEST	RATE
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m <sup>2</sup> Spray with at least 400 L water per hectare to ensure even coverage.
	Larvae of Billbug	Preferably spray on to wet or dewy grass. Irrigate with 12 mm of water commencing within one hour of application.
Roses	Aphids	25 mL/100 L
Ornamental plants	Aphids Azalea lace bug Bronze orange bug Harlequin bug Citrus mealybug Greenhouse thrips Fullers rose weevil	
	Hibiscus flower beetle	50 mL/100 L
	Longtailed mealybug	50 mL/100 L + surfactant
	Psyllids	25 mL/100 L
	Soft scales	25 mL/100 L
Duboisia	Green peach aphid	25 mL/100 L
Pandanus trees	Flatid ( <i>Jamella australiae</i> )	<b>Spot spray</b> 875 mL/100 L of water <b>Stem injection</b> 1.75 L/1 L of water



**CRITICAL COMMENTS**

Apply at peak egg hatch, which is mid spring to mid summer depending on species.

Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.

Apply as a thorough cover spray at first sign of insect infestation.

Spray buds and flowers as needed.

Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.

Spray at first sign and then a week later.

Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.

Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.

**Spot spray:** Spray 100 mL of mixture directly into the leafy throat of each head.

**Stem injection:** Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Pride Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

## SOIL DRENCHED APPLICATIONS

CROP	PEST	RATE
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water
Roses	Aphids	3.5 mL/2 L water/plant



**CRITICAL COMMENTS**

Use as a soil drench application: Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400 to 500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 to 10 days after treatment. Treat at least 6 to 10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.

Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. Do not dilute to the point where mix runs out the bottom of pots.

Use as a soil drench for pots up to 20 L capacity.

Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.

Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Pride Insecticide to the 2 L of water.

For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the Pride Insecticide mixture, and then water it in well immediately after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER,  
CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER  
APPROPRIATE LEGISLATION**

## GENERAL INSTRUCTIONS

GROUP	<b>4A</b>	INSECTICIDE
-------	-----------	-------------

### **Insecticide Resistance Warning**

For insecticide resistance management, Pride Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Pride Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Pride Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Pride Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Amgrow accepts no liability for any losses that may result from the failure of Pride Insecticide to control resistant insects. Pride Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Amgrow representative or local agricultural department agronomist.

**Insecticide Resistance Management Strategy for Aphids other than Green Peach Aphid and Cotton Aphid, Whitefly and Melon Thrips:** DO NOT apply Pride Insecticide in consecutive sprays. Rotate with registered pesticides from other groups.

### **Confined environments such as glasshouses**

Annuals: Do not apply more than one spray of Pride Insecticide to any one crop.

Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

### **Note on Ornamentals**

Pride Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.



### **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Pride Insecticide to water in the spray vat while stirring or with agitators in motion. Pride Insecticide should not be mixed with other pesticides or fertilisers.

### **PROTECTION OF LIVESTOCK**

Dangerous to bees.

DO NOT spray any plants in flower while bees are foraging.

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse container before disposal.

Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

“Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow’s skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions.”

® Trademark of Ecofertiliser

APVMA Approval No. 64900/1LL/0410





# CAUTION

KEEP OUT OF THE REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Pride

Insecticide

ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP

4A

INSECTICIDE

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT:**

**READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**

Amgrow Pty Ltd (Commercial Products Division)

ABN 22 069 900 456

Unit B2A, 3-29 Birnie Ave,

Lidcombe NSW 2141

TEL: 1800 631 008

[www.ecofertiliser.com.au](http://www.ecofertiliser.com.au)

**CONTENTS**

# 1L

® Pride is a registered trademark of Ecofertiliser Pty Ltd.

APVMA Approval No. 64900/1L/0410



### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

### **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

### **CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

### **IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

B/N

DOM



**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

▲ PEEL HERE ▲

**Pride**  
Insecticide



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP **4A** INSECTICIDE

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT:**  
**READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**

Amgrow Pty Ltd (Commercial Products Division)  
ABN 22 069 900 456  
Unit B2A, 3-29 Birnie Ave,  
Lidcombe NSW 2141  
TEL: 1800 631 008  
[www.ecofertiliser.com.au](http://www.ecofertiliser.com.au)

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
APVMA Approval No. 64900/2.5L/0410



File Name: 64900\_49555\_2.5L\_label\_MPL\_V2.pdf

Dimensions: 140mm wide x 130mm high

Scale of Print: A4 100% A3 100%

## DIRECTIONS FOR USE

CROP	PEST	RATE
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m <sup>2</sup> Spray with at least 400 L water per hectare to ensure even coverage. Preferably spray on to wet or dewy grass. Irrigate with 12 mm of water commencing within one hour of application.
	Larvae of Billbug	
Roses	Aphids	25 mL/100 L
Ornamental plants	Aphids	
	Azalea lace bug	
	Bronze orange bug	
	Harlequin bug	
	Citrus mealybug	
	Greenhouse thrips	
Fullers rose weevil		
	Hibiscus flower beetle	50 mL/100 L
	Longtailed mealybug	50 mL/100 L + surfactant
	Psyllids	25 mL/100 L
	Soft scales	25 mL/100 L
Duboisia	Green peach aphid	25 mL/100 L
Pandanus trees	Flatid (Jamella australiae)	<b>Spot spray</b> 875 mL/100 L of water <b>Stem injection</b> 1.75 L/1 L of water



**CRITICAL COMMENTS**

Apply at peak egg hatch, which is mid spring to mid summer depending on species.

Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.

Apply as a thorough cover spray at first sign of insect infestation.

Spray buds and flowers as needed.

Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.

Spray at first sign and then a week later.

Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.

Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.

**Spot spray:** Spray 100 mL of mixture directly into the leafy throat of each head.

**Stem injection:** Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Pride Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

## SOIL DRENCHED APPLICATIONS

CROP	PEST	RATE
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water
Roses	Aphids	3.5 mL/2 L water/plant



**CRITICAL COMMENTS**

Use as a soil drench application: Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400 to 500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 to 10 days after treatment. Treat at least 6 to 10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.

Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. Do not dilute to the point where mix runs out the bottom of pots.

Use as a soil drench for pots up to 20 L capacity.

Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.

Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Pride Insecticide to the 2 L of water.

For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the Pride Insecticide mixture, and then water it in well immediately after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## GENERAL INSTRUCTIONS

GROUP

4A

INSECTICIDE

### **Insecticide Resistance Warning**

For insecticide resistance management, Pride Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Pride Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Pride Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Pride Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Amgrow accepts no liability for any losses that may result from the failure of Pride Insecticide to control resistant insects. Pride Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Amgrow representative or local agricultural department agronomist.

**Insecticide Resistance Management Strategy for Aphids other than Green Peach Aphid and Cotton Aphid, Whitefly and Melon Thrips:** DO NOT apply Pride Insecticide in consecutive sprays. Rotate with registered pesticides from other groups.

### **Confined environments such as glasshouses**

Annuals: Do not apply more than one spray of Pride Insecticide to any one crop.

Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

### **Note on Ornamentals**

Pride Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.





## **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Pride Insecticide to water in the spray vat while stirring or with agitators in motion. Pride Insecticide should not be mixed with other pesticides or fertilisers.

## **PROTECTION OF LIVESTOCK**

Dangerous to bees.

DO NOT spray any plants in flower while bees are foraging.

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

® Trademark of Ecofertiliser

APVMA Approval No. 64900/2.5LL/0410



**CAUTION**  
 KEEP OUT OF THE REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP	<b>4A</b>	INSECTICIDE
-------	-----------	-------------

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT:  
 READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**  
 Amgrow Pty Ltd (Commercial Products Division)  
 ABN 22 069 900 456  
 Unit B2A, 3-29 Birnie Ave,  
 Lidcombe NSW 2141  
 TEL: 1800 631 008  
 www.ecofertiliser.com.au

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
 APVMA Approval No. 64900/2.5L/0410



Page 1 130mm x 116mm

WHILE EVERY EFFORT HAS BEEN MADE TO ENSURE THAT THERE ARE NO ERRORS, IT REMAINS THE CLIENTS RESPONSIBILITY TO PROVIDE THE FINAL INDEPENDANT INSPECTION. PLEASE CHECK THE MATERIAL CAREFULLY.  
 • Please check all copy, legal standards and sign and date your approval for print • Colours are representational only.

SIZE: F: 130 x 116mm

FILE NAME: ECO FERTILISER 2.5L

ART MEMO:

ROLL DIRECTION:

ARTIST: CHRIS

COLOURS: C M Y K



PH (02) 9550 0999 FAX (02) 9550 9734

VERSION:	1. Supplied art	2. Alt art	2. Alt art	2. Alt art				
DATE:	24/05/10	27/05/10	27/05/10	08/06/10				
ALTERATIONS:		Remove Barcode on back	Add code on last page of leaflet	Correction on last page of leaflet				

CLIENT APPROVAL:

DATE:

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

**IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

B/N

DOM



BACK 130mm x 140mm

File Name:  
64900\_49555\_5L\_label\_MPL\_V2.pdf

Dimensions: 140mm wide x 130mm high

Scale of Print: A4 100% A3 100%

## CAUTION

KEEP OUT OF THE REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

▲ PEEL HERE ▲

 **Pride**  
Insecticide



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP	<b>4A</b>	INSECTICIDE
-------	-----------	-------------

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT:**  
**READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**

Amgrow Pty Ltd (Commercial Products Division)  
ABN 22 069 900 456  
Unit B2A, 3-29 Birnie Ave,  
Lidcombe NSW 2141  
TEL: 1800 631 008  
[www.ecofertiliser.com.au](http://www.ecofertiliser.com.au)

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
APVMA Approval No. 64900/5L/0410



CONTENTS  
**5L**

## DIRECTIONS FOR USE

CROP	PEST	RATE
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m <sup>2</sup> Spray with at least 400 L water per hectare to ensure even coverage. Preferably spray on to wet or dewy grass. Irrigate with 12 mm of water commencing within one hour of application.
	Larvae of Billbug	
Roses	Aphids	25 mL/100 L
Ornamental plants	Aphids	
	Azalea lace bug	
	Bronze orange bug	
	Harlequin bug	
	Citrus mealybug	
	Greenhouse thrips	
Fullers rose weevil		
	Hibiscus flower beetle	50 mL/100 L
	Longtailed mealybug	50 mL/100 L + surfactant
	Psyllids	25 mL/100 L
	Soft scales	25 mL/100 L
Duboisia	Green peach aphid	25 mL/100 L
Pandanus trees	Flatid (Jamella australiae)	<b>Spot spray</b> 875 mL/100 L of water <b>Stem injection</b> 1.75 L/1 L of water



**CRITICAL COMMENTS**

Apply at peak egg hatch, which is mid spring to mid summer depending on species.

Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.

Apply as a thorough cover spray at first sign of insect infestation.

Spray buds and flowers as needed.

Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.

Spray at first sign and then a week later.

Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.

Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.

**Spot spray:** Spray 100 mL of mixture directly into the leafy throat of each head.

**Stem injection:** Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Pride Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

## SOIL DRENCHED APPLICATIONS

CROP	PEST	RATE
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water
Roses	Aphids	3.5 mL/2 L water/plant





**CRITICAL COMMENTS**

Use as a soil drench application: Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400 to 500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 to 10 days after treatment. Treat at least 6 to 10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.

Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. Do not dilute to the point where mix runs out the bottom of pots.

Use as a soil drench for pots up to 20 L capacity.

Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.

Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Pride Insecticide to the 2 L of water.

For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the Pride Insecticide mixture, and then water it in well immediately after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## GENERAL INSTRUCTIONS

GROUP

4A

INSECTICIDE

### **Insecticide Resistance Warning**

For insecticide resistance management, Pride Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Pride Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Pride Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Pride Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Amgrow accepts no liability for any losses that may result from the failure of Pride Insecticide to control resistant insects. Pride Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Amgrow representative or local agricultural department agronomist.

**Insecticide Resistance Management Strategy for Aphids other than Green Peach Aphid and Cotton Aphid, Whitefly and Melon Thrips:** DO NOT apply Pride Insecticide in consecutive sprays. Rotate with registered pesticides from other groups.

### **Confined environments such as glasshouses**

Annuals: Do not apply more than one spray of Pride Insecticide to any one crop.

Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

### **Note on Ornamentals**

Pride Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.



## **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Pride Insecticide to water in the spray vat while stirring or with agitators in motion. Pride Insecticide should not be mixed with other pesticides or fertilisers.

## **PROTECTION OF LIVESTOCK**

Dangerous to bees.

DO NOT spray any plants in flower while bees are foraging.

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

® Trademark of Ecofertiliser

APVMA Approval No. 64900/5LL/0410



**CAUTION**  
 KEEP OUT OF THE REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP	<b>4A</b>	INSECTICIDE
-------	-----------	-------------

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT:**  
 READ THE ATTACHED LEAFLET BEFORE USE

**Distributed by:**  
 Amgrow Pty Ltd (Commercial Products Division)  
 ABN 22 069 900 456  
 Unit B2A, 3-29 Birnie Ave,  
 Lidcombe NSW 2141  
 TEL: 1800 631 008  
 www.ecofertiliser.com.au

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
 APVMA Approval No. 64900/5L/0410



PAGE 1 130mm x 116mm

WHILE EVERY EFFORT HAS BEEN MADE TO ENSURE THAT THERE ARE NO ERRORS, IT REMAINS THE CLIENTS RESPONSIBILITY TO PROVIDE THE FINAL INDEPENDANT INSPECTION. PLEASE CHECK THE MATERIAL CAREFULLY.  
 • Please check all copy, legal standards and sign and date your approval for print • Colours are representational only.

SIZE: F: 130 x 116mm

FILE NAME: ECO FERTILISER 5L

ART MEMO:

ROLL DIRECTION:

ARTIST: CHRIS

COLOURS: C M Y K



PH (02) 9550 0999 FAX (02) 9550 9734

VERSION:	1. Supplied art	2. Alt art	3. Alt art	4. Alt art				
DATE:	24/05/10	27/05/10	27/05/10	08/06/10				
ALTERATIONS:		Remove Barcode on back	Add code on last page of leaflet	Correction on last page of leaflet				

CLIENT APPROVAL:

DATE:

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

**IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

B/N

DOM



BACK 130mm x 140mm

File Name:  
64900\_49555\_10L\_label\_MPL\_V2.pdf

Dimensions: 220mm wide x 170mm high

Scale of Print: A4 100%    A3 100%

**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PEEL HERE



# Pride

Insecticide



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

**GROUP 4A INSECTICIDE**

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**  
Amgrow Pty Ltd (Commercial Products Division)  
ABN 22 069 900 456  
Unit B2A, 3-29 Birnie Ave,  
Lidcombe NSW 2141  
TEL: 1800 631 008  
[www.ecofertiliser.com.au](http://www.ecofertiliser.com.au)

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
APVMA Approval No. 64900/10L/0410



**CONTENTS**  
**10L**

## DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m <sup>2</sup> Spray with at least 400 L water per hectare to ensure even coverage. Preferably spray on to wet or dewy grass.	Apply at peak egg hatch, which is mid spring to mid summer depending on species.
	Larvae of Billbug	Irrigate with 12 mm of water commencing within one hour of application.	Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.
Roses	Aphids	25 mL/100 L	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids Azalea lace bug Bronze orange bug Harlequin bug Citrus mealybug Greenhouse thrips Fullers rose weevil		
	Hibiscus flower beetle	50 mL/100 L	Spray buds and flowers as needed.
	Longtailed mealybug	50 mL/100 L + surfactant	Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	25 mL/100 L	Spray at first sign and then a week later.
	Soft scales	25 mL/100 L	Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid	25 mL/100 L	Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus trees	Flatid ( <i>Jamella australiae</i> )	Spot spray 875 mL/100 L of water Stem injection 1.75 L/1 L of water	Spot spray: Spray 100 mL of mixture directly into the leafy throat of each head. Stem injection: Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Pride Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.





## SOIL DRENCHED APPLICATIONS

CROP	PEST	RATE	CRITICAL COMMENTS
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height	Use as a soil drench application: Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400 to 500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 to 10 days after treatment. Treat at least 6 to 10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant	Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. Do not dilute to the point where mix runs out the bottom of pots.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot	Use as a soil drench for pots up to 20 L capacity.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water	Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.
Roses	Aphids	3.5 mL/2 L water/plant	Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Pride Insecticide to the 2 L of water.

For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the Pride Insecticide mixture, and then water it in well immediately after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**



## GENERAL INSTRUCTIONS

GROUP **4A** INSECTICIDE

### **Insecticide Resistance Warning**

For insecticide resistance management, Pride Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Pride Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Pride Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Pride Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Amgrow accepts no liability for any losses that may result from the failure of Pride Insecticide to control resistant insects. Pride Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Amgrow representative or local agricultural department agronomist.

### **Insecticide Resistance Management Strategy for Aphids other than Green Peach Aphid and Cotton Aphid,**

**Whitefly and Melon Thrips:** DO NOT apply Pride Insecticide in consecutive sprays. Rotate with registered pesticides from other groups.

### **Confined environments such as glasshouses**

Annuals: Do not apply more than one spray of Pride Insecticide to any one crop.

Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

### **Note on Ornamentals**

Pride Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

### **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Pride Insecticide to water in the spray vat while stirring or with agitators in motion. Pride Insecticide should not be mixed with other pesticides or fertilisers.

### **PROTECTION OF LIVESTOCK**

Dangerous to bees.

DO NOT spray any plants in flower while bees are foraging.

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.



## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

## **CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

® Trademark of Ecofertiliser

APVMA Approval No. 64900/10LL/0410



**CAUTION**  
 KEEP OUT OF THE REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

**GROUP 4A INSECTICIDE**

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**

Amgrow Pty Ltd (Commercial Products Division)  
 ABN 22 069 900 456  
 Unit B2A, 3-29 Birnie Ave,  
 Lidcombe NSW 2141  
 TEL: 1800 631 008  
 www.ecofertiliser.com.au

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
 APVMA Approval No. 64900/10L/0410



Page 1 196mm x 170mm

WHILE EVERY EFFORT HAS BEEN MADE TO ENSURE THAT THERE ARE NO ERRORS, IT REMAINS THE CLIENTS RESPONSIBILITY TO PROVIDE THE FINAL INDEPENDANT INSPECTION. PLEASE CHECK THE MATERIAL CAREFULLY.  
 \*Please check all copy, legal standards and sign and date your approval for print \* Colours are representational only.

SIZE: F: 170 x 196 mm FILE NAME: ECO FERTILISER 10L  
 BASE SIZE: F: 170 x 220 mm ROLL DIRECTION:

ART MEMO:  
 ARTIST: CHRIS

COLOURS: C M Y K



PH (02) 9550 0999 FAX (02) 9550 9734

VERSION:	1. Supplied art	2. Alt art	3. Alt art	4. Alt art				
DATE:	26/05/10	27/05/10	27/05/10	08/06/10				
ALTERATIONS:		Remove Barcode on back	Add code on last page of leaflet	corrections on last page of leaflet and page 3				

CLIENT APPROVAL:

DATE:

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

**IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

B/N

DOM0



File Name:  
64900\_49555\_20L\_label\_MPL\_V2.pdf

Dimensions: 220mm wide x 170mm high

Scale of Print: A4 100%  
A3 100%

**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PEEL HERE

 **Pride**  
Insecticide



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP **4A** INSECTICIDE

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**  
Amgrow Pty Ltd (Commercial Products Division)  
ABN 22 069 900 456  
Unit B2A, 3-29 Birnie Ave,  
Lidcombe NSW 2141  
TEL: 1800 631 008  
[www.ecofertiliser.com.au](http://www.ecofertiliser.com.au)

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
APVMA Approval No. 64900/20L/0410



**CONTENTS**  
**20L**

## DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
Turf	First instar larvae of: African black beetle Argentinian scarab Pruinose scarab	2.5 L/ha or 25 mL/100 m <sup>2</sup> Spray with at least 400 L water per hectare to ensure even coverage. Preferably spray on to wet or dewy grass.	Apply at peak egg hatch, which is mid spring to mid summer depending on species.
	Larvae of Billbug	Irrigate with 12 mm of water commencing within one hour of application.	Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.
Roses	Aphids	25 mL/100 L	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids Azalea lace bug Bronze orange bug Harlequin bug Citrus mealybug Greenhouse thrips Fullers rose weevil		
	Hibiscus flower beetle	50 mL/100 L	Spray buds and flowers as needed.
	Longtailed mealybug	50 mL/100 L + surfactant	Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	25 mL/100 L	Spray at first sign and then a week later.
	Soft scales	25 mL/100 L	Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid	25 mL/100 L	Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus trees	Flatid ( <i>Jamella australiae</i> )	Spot spray 875 mL/100 L of water Stem injection 1.75 L/1 L of water	Spot spray: Spray 100 mL of mixture directly into the leafy throat of each head. Stem injection: Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Pride Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.



## SOIL DRENCHED APPLICATIONS

CROP	PEST	RATE	CRITICAL COMMENTS
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height	Use as a soil drench application: Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400 to 500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 to 10 days after treatment. Treat at least 6 to 10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Seedling eucalypts (to 1 m high) in pots	Chrysomelid beetle larvae Psyllids	2.5 mL/plant	Mix in water up to 0.5 L per 3 L pot and apply to soil. Use less water for smaller pots. Do not dilute to the point where mix runs out the bottom of pots.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot	Use as a soil drench for pots up to 20 L capacity.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water	Use as a soil drench. 5 L of mixture will treat twenty 6 L pots.
Roses	Aphids	3.5 mL/2 L water/plant	Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Pride Insecticide to the 2 L of water.

For soil drench treatments, remove mulch and dead vegetation, and moisten the soil surface first. Apply the Pride Insecticide mixture, and then water it in well immediately after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**





## GENERAL INSTRUCTIONS

GROUP **4A** INSECTICIDE

### **Insecticide Resistance Warning**

For insecticide resistance management, Pride Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Pride Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Pride Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Pride Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Amgrow accepts no liability for any losses that may result from the failure of Pride Insecticide to control resistant insects. Pride Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Amgrow representative or local agricultural department agronomist.

### **Insecticide Resistance Management Strategy for Aphids other than Green Peach Aphid and Cotton Aphid,**

**Whitefly and Melon Thrips:** DO NOT apply Pride Insecticide in consecutive sprays. Rotate with registered pesticides from other groups.

### **Confined environments such as glasshouses**

Annuals: Do not apply more than one spray of Pride Insecticide to any one crop.

Perennials: Rotate with registered insecticides from other groups. Use a maximum of three sprays in any 12 month period.

### **Note on Ornamentals**

Pride Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

### **MIXING**

Prior to pouring, shake container vigorously, then add the required quantity of Pride Insecticide to water in the spray vat while stirring or with agitators in motion. Pride Insecticide should not be mixed with other pesticides or fertilisers.

### **PROTECTION OF LIVESTOCK**

Dangerous to bees.

DO NOT spray any plants in flower while bees are foraging.

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.



## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS: Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

## **CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

® Trademark of Ecofertiliser

APVMA Approval No. 64900/20LL/0410



**CAUTION**  
 KEEP OUT OF THE REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

**GROUP 4A INSECTICIDE**

For the control of a range of insect pests in turf and ornamental situations as per the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**Distributed by:**

Amgrow Pty Ltd (Commercial Products Division)  
 ABN 22 069 900 456  
 Unit B2A, 3-29 Birnie Ave,  
 Lidcombe NSW 2141  
 TEL: 1800 631 008  
 www.ecofertiliser.com.au

© Pride is a registered trademark of Ecofertiliser Pty Ltd.  
 APVMA Approval No. 64900/20L/0410



Page 1 196mm x 170mm

WHILE EVERY EFFORT HAS BEEN MADE TO ENSURE THAT THERE ARE NO ERRORS, IT REMAINS THE CLIENTS RESPONSIBILITY TO PROVIDE THE FINAL INDEPENDANT INSPECTION. PLEASE CHECK THE MATERIAL CAREFULLY.  
 \*Please check all copy, legal standards and sign and date your approval for print \* Colours are representational only.

SIZE: F: 170 x 196 mm	FILE NAME: ECO FERTILISER 20L	ART MEMO:	
	ROLL DIRECTION:	ARTIST: CHRIS	
COLOURS: <span style="background-color: cyan; border: 1px solid black; padding: 2px;">C</span> <span style="background-color: magenta; border: 1px solid black; padding: 2px;">M</span> <span style="background-color: yellow; border: 1px solid black; padding: 2px;">Y</span> <span style="background-color: black; border: 1px solid black; padding: 2px;">K</span>			
	PH (02) 9550 0999 FAX (02) 9550 9734		

VERSION:	1. Supplied art	2. Alt art	3. Alt art	4. Alt art				
DATE:	26/05/10	27/05/10	27/05/10	08/06/10				
ALTERATIONS:		Remove Barcode on back	Add code on last page of leaflet	corrections on last page of leaflet and page 3				

CLIENT APPROVAL: \_\_\_\_\_ DATE: \_\_\_\_\_

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

**CONDITIONS OF SALE**

"Amgrow Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Amgrow's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Amgrow has any authority to add to or alter these conditions."

**IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

B/N

DOM0

