



**Company Name:** BAYER CROPSCIENCE PTY LTD  
**Product Name:** RESERVE STRESSGARD TURFFUNGICIDE  
**APVMA Approval No:** 81269/105929

<b>Label Name:</b>	RESERVE STRESSGARD TURF FUNGICIDE
--------------------	-----------------------------------

<b>Signal Headings:</b>	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	---

<b>Constituent Statements:</b>	720 g/L CHLOROTHALONIL
--------------------------------	------------------------

<b>Mode of Action:</b>	<table border="1"><tr><td>GROUP</td><td><b>M5</b></td><td>FUNGICIDE</td></tr></table>	GROUP	<b>M5</b>	FUNGICIDE
GROUP	<b>M5</b>	FUNGICIDE		

<b>Statement of Claims:</b>	For the control of fungal diseases in turf.
-----------------------------	---

<b>Net Contents:</b>	5L-1000L
----------------------	----------

<b>Restraints:</b>	DO NOT mix spraying oils or wetting agents with this product.
--------------------	---

<b>Directions for Use:</b>	This section contains file attachment. <b>File Name:</b> RESERVE STRESSGARD TURF FUNGICIDE Directions for Use table 2016.03.07.pdf <b>File Size:</b> 65175 bytes
----------------------------	--

<b>Other Limitations:</b>	
---------------------------	--

<b>Withholding Periods:</b>	
-----------------------------	--

<b>Trade Advice:</b>	
----------------------	--

<b>General Instructions:</b>	<p>RESERVE STRESSGARD is a protectant fungicide for the control of fungal diseases in turf. For effective disease control a good leaf cover must be achieved before a widespread disease outbreak occurs.</p> <p><b>MIXING</b> Thoroughly shake and invert the pack several times before opening. Half-fill spray tank and with agitation system running, add the required amount of product.</p> <p><b>APPLICATION RATES PER HECTARE</b> Low volume: <math>\geq 200</math> L/ha High volume: <math>\geq 500</math> L/ha The volume of water applied should increase as the plants grow in size to maintain adequate cover.</p> <p><b>COMPATIBILITY</b> This product is compatible with wettable powder and suspension concentrate formulations of most commonly used fungicides, insecticides and miticides. Unless experience in your area has shown other combinations to be physically compatible and non-injurious to your crop, do not combine with other chemicals, especially oil-based emulsifiable or suspension concentrate pesticides. This product should not be mixed with spraying oils or sprayed onto plants that have been sprayed with oil for at least 10 days after the last spray.</p> <p><b>WETTING AGENTS</b> The addition of surfactants does not improve this product's efficacy and under some conditions may cause plant injury.</p>
------------------------------	--

<b>Resistance Warning:</b>	<p>RESERVE STRESSGARD is a member of the multi-site activity group of fungicides. For fungicide resistance management, this product is a Group M5 fungicide. Some naturally occurring individual fungi resistant to this product and other Group M5 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group M5 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.</p>
----------------------------	--

<b>Precautions:</b>	
---------------------	--

<b>Protections:</b>	<p><b>PROTECTION OF LIVESTOCK</b> DO NOT contaminate feed or foodstuffs with this product or empty containers.</p> <p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b> Dangerous to fish. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.</p>
---------------------	--

<b>Storage and Disposal:</b>	<p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose,</p>
------------------------------	--

clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. Do not re-use empty containers for any other purpose.

**Safety Directions:**

Attacks eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length pvc gloves and goggles and disposable mist face mask covering mouth and nose. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

**First Aid Instructions:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do not induce vomiting. Give a glass of water.

**First Aid Warnings:**

## RESERVE STRESSGARD TURF FUNGICIDE

### DIRECTIONS FOR USE

SITUATION	DISEASE	STATE	RATE	CRITICAL COMMENTS
Turf and lawns, bowling greens, golf greens	Dollar spot ( <i>Sclerotinia homoeocarpa</i> )	Qld, NSW, ACT, Vic, SA, WA only	130 – 200 mL/100 m <sup>2</sup> in 20-60 L water	If disease is already established, use the higher rate. Apply at 7 – 14-day intervals. Under severe conditions, spray at 7-day intervals. Do not mow or water treated area until turf or lawn is thoroughly dry.
	Brown patch ( <i>Rhizoctonia</i> sp.)	All states		
	Gray leaf spot ( <i>Pyricularia grisea</i> )	NSW, WA only	24 L/ha	Apply to leaf as a curative application. Do not irrigate for 24 hours after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**